Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DEATHS FROM INFLUENZA AND PNEUMONIA COMBINED.

COMPARISON OF THE FIRST SEVEN WEEKS OF THE YEARS 1919–1922, INCLUSIVE, FOR CERTAIN LARGE CITIES OF THE UNITED STATES.

The accompanying table gives the number of reported deaths from influenza and pneumonia (all forms), combined, during the first seven weeks of the years 1919, 1920, 1921, and 1922, in 36 large cities of the United States.

This is a continuation of the table printed on pages 333–334 of the Public Health Reports of February 17, 1922 (vol. 37, No. 7).

The weeks for which figures are given all ended on Saturday, the "first" week for each year ending on the following days, respectively: January 4, 1919, January 10, 1920, January 8, 1921, and January 7, 1922.

The figures for 1919 and 1920 were taken from the Weekly Health Index, issued by the Bureau of the Census, Department of Commerce, supplemented by reports to the Public Health Service. For 1921 and 1922 the figures are taken from reports made by the city health officers to the Public Health Service.

Blanks in the table indicate that no reports of deaths from influenza or pneumonia were received for the week. This does not always indicate that no deaths from these diseases occurred. In the seventh week of 1922 it means in most instances that the report has been delayed.

Number of deaths from influenza and pneumonia (all forms) combined.

City.	Week number.						
	First.	Second.	Third.	Fourth.	Fifth.	Sixth.	Seventh.
Birmingham, Ala.:							
1922	8	10	14	6	13	4	4
1921	7	14	6	4	9	9	12
1920	13	9	16	14	22	18	59
1919	36	44	52	41	29	21	28
Los Angeles, Calif.:							
1922	18	19	14	21	26	29	33
1921	12	19	9	13	15	12	17
1920	16	18	19	22	42	88	74
1919	99	151	178	177	104	47	21
Oakland, Calif.:							
1922	4	5	5	6	8	8	12
1921	4	3	8	7	9	4	6
1920	4	8	20	24	55	54	60
1919	66	92	111	67	38	18	[1] 8
San Francisco, Calif.:							
1922	11	12	4	12	9	15	36
1921	.....	5	8	9	7	11	13
1920	14	26	48	59	115	137	113
1919	194	290	310	149	59	41	20
Denver, Colo.:							
1922	22	11	10	17	18	16	19
1921	25	22	23	11	16	21	20
1920	21	18	24	49	159	160	67
1919	65	47	35	24	29	30	37
New Haven, Conn.:							
1922	5	1	5	4	13	10	14
1921	4	7	7	7	2	6	9
1920	6	8	10	19	20	60	68
1919	40	38	27	26	20	12	11

[1] Pneumonia (all forms) deaths only.

Number of deaths from influenza and pneumonia (all forms) combined—Continued.

City.	Week number.						
	First.	Second.	Third.	Fourth.	Fifth.	Sixth.	Seventh.
Washington, D. C.:							
1922	20	22	27	27	25	22	27
1921	22	22	14	9	9	12	19
1920	22	27	81	181	164	92	55
1919	139	109	107	73	60	42	40
Atlanta, Ga.:							
1922	13	7	9	7	20	17	11
1921	10	8	9	5	7	18	10
1920	19	11	10	15	32	75	104
1919	[1] 40	[1] 40	[1] 54	[1] 57	[1] 54	[1] 28	[1] 21
Chicago, Ill.:							
1922	48	43	63	65	72	80	56
1921	64	79	89	102	92	90	75
1920	107	153	472	1,109	1,005	494	243
1919	321	269	328	341	277	194	235
Indianapolis, Ind.:							
1922	20	11	9	17	29	42	39
1921	15	12	13	13	21	6	13
1920	18	16	21	36	92	124	72
1919	34	40	25	28	25	23	28
Louisville, Ky.:							
1922	6	12	18	7	16	24	28
1921	6	4	5	5	2	2	9
1920	10	10	9	18	40	52	48
1919	22	20	21	30	20	19	19
New Orleans, La.:							
1922	13	14	14	13	4	25	20
1921	18	18	21	[2] 3	12	21	23
1920	27	27	27	32	36	62	89
1919	94	141	202	201	125	58	49
Baltimore, Md.:							
1922	32	25	24	26	29	27	29
1921	33	20	24	18	26	56	44
1920	20	35	24	59	122	268	231
1919	48	75	83	150	138	126	117
Boston, Mass.:							
1922	21	17	36	28	33	38	51
1921	27	23	36	33	22	10	26
1920	28	28	45	85	158	255	216
1919	244	227	158	153	110	89	71
Cambridge, Mass.:							
1922	5	8	3	4	7	7	8
1921	4	5	5	5	1	3	4
1920	8	7	8	14	22	28	23
1919	39	22	20	16	[2] 5	10	3
Fall River, Mass.:							
1922	5	4	3	6	5	7	9
1921	14	5	11	4	5	8	5
1920	7	10	5	3	5	16	25
1919	10	18	16	14	17	17	15
Lowell, Mass.:							
1922	4	7	5	4	4	6	5
1921	7	6	8	3	6	4	2
1920	5	4	2	7	12	10	36
1919	13	[1] 10	20	26	11	[1] 7	18
Worcester, Mass.:							
1922	5	10	11	7	16	16	16
1921	4	7	13	9	4	10	12
1920	10	9	7	14	15	44	52
1919	40	36	44	22	23	21	23
Minneapolis, Minn.:							
1922	10	6	9	9	6	9	4
1921	13	14	10	8	10	16	20
1920	12	10	9	63	168	125	53
1919	37	45	24	32	31	31	14
St. Paul, Minn.:							
1922	7	13	7	3	8	6	6
1921	9	5	9	9	.....	7	8
1920	4	10	26	75	80	63	26
1919	39	25	14	12	15	13	11
Kansas City, Mo.:							
1922	15	13	14	25	25	28	39
1921	17	17	19	13	14	17	16
1920	13	29	96	120	220	167	74
1919	49	50	68	45	58	40	51
Omaha, Nebr.:							
1922	11	9	17	12	16	12	11
1921	8	7	4	14	.....	4	12
1920	4	7	13	45	62	63	32
1919	25	25	17	17	11	12	10

[1] Pneumonia (all forms) deaths only. [2] Influenza deaths only.

Number of deaths from influenza and pneumonia (all forms) combined—Continued.

City.	Week number.						
	First.	Second.	Third.	Fourth.	Fifth.	Sixth.	Seventh.
Newark, N. J.:							
1922	13	15	20	20	33	29	39
1921	18	14	15	7	12	13	12
1920	17	14	30	55	116	142	93
1919	72	66	57	53	50	45	32
Buffalo, N. Y.:							
1922	6	20	13	19	21	15	15
1921	20	18	18	20	13	18	20
1920	10	7	19	17	67	141	145
1919	48	[1] 19	90	123	90	75	35
New York, N. Y.:							
1922	215	263	284	302	481	596	576
1921	235	216	204	203	199	212	212
1920	218	261	511	1,308	1,988	1,796	987
1919	753	870	998	1,193	1,153	893	786
Rochester, N. Y.:							
1922	5	11	12	14	6	7	14
1921	4	3	6	8	5	5	4
1920	13	7	12	23	50	52	27
1919	59	26	17	21	12	16	16
Syracuse, N. Y.:							
1922	4	6	4	6	7	7	.....
1921	4	8	3	5	6	2	7
1920	9	8	10	31	89	78	29
1919	8	13	4	14	18	10	10
Cincinnati, Ohio:							
1922	14	20	15	19	21	27	41
1921	14	16	13	11	18	16	17
1920	14	12	17	25	38	62	81
1919	51	18	18	26	23	39	37
Cleveland, Ohio:							
1922	.....	.....	30	28	25	18	25
1921	25	22	23	24	31	28	31
1920	21	25	26	41	158	258	177
1919	132	94	92	92	108	100	80
Columbus, Ohio:							
1922	5	9	4	10	8	6	10
1921	8	8	12	12	13	12	7
1920	15	9	8	22	59	118	66
1919	15	14	10	20	19	11	15
Toledo, Ohio:							
1922	6	9	8	12	7	6	5
1921	.....	3	9	10	5	4	3
1920	9	8	9	18	54	50	50
1919	19	15	19	20	15	6	11
Portland, Oreg.:							
1922	4	7	4	6	5	15	17
1921	6	5	7	6	4	8	5
1920	13	8	9	17	21	57	52
1919	55	101	123	122	50	15	10
Philadelphia, Pa.:							
1922	73	98	87	86	85	91	101
1921	72	83	85	101	114	108	115
1920	55	75	108	153	289	564	620
1919	142	194	229	259	308	262	232
Providence, R. I.:							
1922	13	8	12	17	11	15	26
1921	14	6	5	8	14	11	9
1920	12	13	8	14	39	88	92
1919	47	59	62	61	35	30	28
Nashville, Tenn.:							
1922	2	7	.....	3	5	5	4
1921	2	8	4	.....	10	9	9
1920	6	11	6	12	8	23	47
1919	20	17	21	21	17	15	16
Richmond, Va.:							
1922	8	9	9	4	8	9	12
1921	5	5	13	6	5	7	10
1920	2	9	6	21	35	38	28
1919	50	26	34	30	23	11	9
Total:							
1922	671	761	823	872	1,125	1,294	1,362
1921	750	737	768	725	738	800	836
1920	802	947	1,771	3,820	5,657	5,922	4,314
1919	3,165	3,346	3,688	3,756	3,180	2,427	2,167

[1] Pneumonia (all forms), deaths only.